

Table 2: Estimates of HIV-2 Seroprevalence

		Capital/Major City				Outside Major City				Urban City Sources		Outside City Sources	
.	Region and Country	Low Risk	.	High Risk	.	Low Risk	.	High Risk	.	Low Risk	High Risk	Low Risk	High Risk
	AFRICA												
*	Benin	0.3	a	-		0.2	a	0.1	a	A0275		A0275	A0336
*	Burkina Faso	0.6	a	4.9	a	0.2	a	-		B0506	O0113	B0506	
*	Cameroon	0		-		0.4		-		C0420		C0420	
*	Congo, Dem. Rep.	-		0		-		-			M0549		
*	Cote d'Ivoire	0.7	a	-		1	ac	-		K0434		K0434	
*	Gambia, The	0.7		-		1.8		-		C0385		C0385	
*	Ghana	0.2	ac	7.1	a	1.4	a	-		N0350	A0300	N0350	
*	Guinea-Bissau	3.8	a	-		9.6		-		N0295		C0265	
*	Mali	-		2.9		-		5.6			M0676		M0676
*	Niger	-		7.1	a	-		8.2	a		H0141		N0333
*	Senegal	0.1		5.2		0.6		7.6		M0773	M0773	M0773	M0773

- No data found.

* See Table 1 for HIV-1 data.

a Rate represents infection with HIV-2 only and dual infection

(HIV-1 and HIV-2), therefore, addition of rates from tables 1 and 2 is not advised.

b Data are best available but are not necessarily reliable due to small (<100) or unknown sample size.

c Data combined.

Notes:

Definitions:

High risk -- prostitutes and clients, STD patients, or other persons with known risk factors.

Low risk -- pregnant women, blood donors, or other persons with no known risk factors.

[Source: U.S. Bureau of the Census, HIV/AIDS Surveillance Data Base.](#)

Internet Release: April 2005